What can YOUth do about Rx Drug Abuse?

Reducing Social Access and Increasing Capacity



Scope of the Problem



- 6.4 Million Americans abused Rx drugs in 2012.
- A majority of them got those drugs from their family or a friend's medicine cabinet.
- Most teens who misuse prescription painkillers are given them for free by a friend or relative.

Lasting Effects



- Recently, prescription painkillers have resulted in more deaths than cocaine and heroin COMBINED.
- Four in five new heroin users started out misusing prescription painkillers.

Local Conditions

Ferry County in 2014

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	4%	U
10 th grade	2%	5%
12 th grade	U	U

Ferry County in 2016

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	U	U
10 th grade	17%	29%
12 th grade	8%	7%

Local Conditions

Ferry County in 2014

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	4%	U
10 th grade	2%	5%
12 th grade	U	U

Ferry County in 2016

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	U	U
10 th grade	17%	29%
12 th grade	8%	7%

Local Conditions

Washington State in 2014

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	2%	4%
10 th grade	5%	8%
12 th grade	6%	9%

Washington State in 2016

	Using Rx drugs to get high in the past 30 days	Using Rx drugs not prescribed to them
8 th grade	2%	5%
10 th grade	4%	8%
12 th grade	5%	9%

Now It's Your Turn!

- Whose support do you need to succeed?
- How are you going to increase community readiness?
- What strategies can you use?
- What resources do you need?



Increasing Community Readiness for Action

- Document and share evidence of problem
- Community
 Conversations about problem behavior







What one group of teens did to tackle the problem

- Applied for a CPWI Enhancement grant
- Researched drug take-back programs
- Developed partnership with Republic City Police
- Earned Republic City Council endorsement





What one group of teens did to tackle the problem

- Launched the Awareness Campaign
- Count It!
- Lock It!
- Drop It!
- Distributed medication lock boxes





Spread the Word

Kick Buns

- Social Media Tools
- Old School Flyers
- Unusual Advertising





Too Much Success

- Be prepared for too much of a good thing!
- Considerations for adoption of a drug take-back program



King County Heroin and Prescription Opiate Task Force – Primary Prevention Workgroup





Opioid Medication & Pain - Flyer

Commonly

Codeine Dilaudid

Fentanyl

prescribed opioids:

Hydrocodone (Vicodin)

Oxymorphone (Opona)

Oxycodone (OxyConfin)

These are only some

of the prescription

optoids. If you get a

prescription for pain, ask your doctor if it is

Hydromorphone

Methodone

Meperidine Morphine

M5 Confin

Percocet

an optoid.





Opioid Medication & Pain: What You Need to Know

If you've had an injury, surgery or major dental work, you are likely to have pain. Pain is a normal part of life and healing. Talk with your doctor about how you can get the most effective pain relief with the least risk.

NON-OPIOID PAIN TREATMENTS HAVE FEWER RISKS

For pain that will likely be gane in a week or two, it is always best to start with nonopicid pain treatments. Opicids may help control pain of lists, but they are usedly not necessary. Consider other costons that may work test as well but how for fawer risks.

- Over-fhe-counter path relievers
- Physical therapy
- Experien
- · Professional help coping with the emotional effects of pain

OPIOIDS ARE STRONG PRESCRIPTION MEDICATIONS

Optoids can be the right choice for treating severe pain, such as from cancer or immediately after major surgery. However, medications such as Vicadin, Percocut and OxyContin are very powerful and can be deadly. Even if you take them as directed, All optoids have serious side effects such as addiction and overdose.

OPIOIDS ARE CHEMICAL COUSINS OF HEROIN AND ARE HIGHLY ADDICTIVE

You can build up a tolerance to optotds over time, so you need to take more and more to get the same relief. The higher the dose, the more dangerous optotds are. You can even become addicted after a short time.

If you are prescribed an opioid for short-term pain:



The prescription should only be for a threeto seven day supply (often this is as few as 10 ptlls).



Take the lowest dose possible for the shortest period of time.



Always talk with the doctor about managing your Dispose of optoid medicines as soon as you stop using them.*

"King County has dozens of pharmacies that can dispose of your snased medications for hee. Find the one closest to you at www.takebackyourneds.org.

www.WashingtonHadthAlliance.org www.BreeCollaborative.org

Opioids Facts & Figures





THE OPIOID EPIDEMIC

Drug overdose is the **leading cause** of accidental death in the US and in Washington state, and opioid addiction is driving the epidemic.

American Society of Addiction Medicine



In 2015 alone, 718 people in Washington died from overdoses involving optoids.

Washington State Department of Health



More than 3 out of 5 drug averdose deaths involve an opioid.

Centers for Disease Control and Prevention



Nationally, 4 in 5 new heroin users started out misusing prescription painkillers.

American Society of Addiction Medicine



In Washington, 57% of people currently using heroin were dependent on prescription opioids before they began using heroin.

UW Alcohol and Drug Abuse Institute

TEENS AND OPIOIDS

1 in 5 teens experiment with prescription drugs.

Centers for Disease Control and Prevention





Nearly half of young people who inject heroin started off abusing prescription drugs.

National Institute on Drug Abuse

More than 4 in 10 teens who misused or abused a prescription drug took it from their parent's medicine cabinat.

Partnership for Drug-Free Kids



95% of parents believe their child has never taken a prescription drug for a reason other than its intended use.

Portnership for Drug-Free Kids

KEEPING KIDS & TEENS SAFE

Sometimes kids and teens are prescribed optaids when they shouldn't be. Or they may be given a prescription for more pills than they need. Teens may also experiment with drugs they find in your medicine cabinet.

Follow these 3 simple guidelines whenever possible:

DON'T FILL a prescription for more than a 3-day supply (maximum 10 pills) for anyone 20 years old or younger. 2 SECURELY STORE optoids away from kids and teens.


3 SAFELY DISPOSE* of extra optoids when you are done.

*King County has dozen all phormacies that can dispose of your unused medications for thes. Find the one closest to you at www.hdusbadkyourneds.org.


www.WashingtonHealthAllianas.org www.BrseCollaborative.org

Find a Medicine Take-Back Location





By List



For questions about Take-back locations in King county, please visit King County Secure MedicineReturn.

Opioid Overdose Reversal - Naloxone



Injectable



Auto-injector



Intranasal



Nasal spray

http://stopoverdose.org/

WA Good Samaritan Law...Means What?

Don't leave a friend to die from a drug overdose.



In Washington State:

If you seek medical help for someone having an overdose, neither of you can be charged for having or using a small amount of drugs.

The Good Samaritan Overdose Law protects you!

CALL 911

*The Good Samaritan Law does not apply to:

- ·outstanding warrants
- probation or parole violations
- drug manufacture or delivery
- ·controlled substances homicide
- other crimes besides drug possession

In these cases, a judge can give you a shorter sentence.

See an overdose

CALL 911!

Washington's **Good Samaritan Law** protects you!

If you get medical help for an overdose or alcohol poisoning, you and the victim cannot be charged for drug use, possession, or underage drinking.

WA RCW 69:50:315

Drug and alcohol overdoses kill at least two people per day in Washington State.

See an overdose **CALL 911!**

Washington's Good Samaritan Law protects you!

Never let someone

"sleep it off."

Learn more at www.stopoverdose.org



http://stopoverdose.org/

See an overdose **CALL 911!**

Washington's Good Samaritan law protects you!



If you get medical help for an overdose or alcohol poisoning, you and the victim cannot be charged for drug use, possession, or underage drinking.

> Never let someone "sleep it off."


WA RCW 69 50 315

Online Resources to Get You Started!

- Stopoverdose.org
- Stopmedicineabuse.org
- Overdoseday.com
- 866teenlink.org
- takebackyourmeds.org
- warecoveryhelpline.org
- O PreventMedAbuse.org













Think Outside the Box...

- InternationalVeterinary Academyof Pain Management
- O https://ivapm.org



Your Workshop Preventionistas!



- O Erin James, MBA, CPP
- Outreach Marijuana & Opiate Prevention Coordinator
- County Behavioral Health and Recovery Division
- O <u>Erin.James@kingcounty.gov</u>
- O 206.477.7657

Your Workshop Preventionistas!

- Janine Koffel, Independent Health Promotion and Prevention Professional
- O Janine.Koffel@gmail.com
- **O** 509-207-9174

